

Choosing AEDs in special situations

Kanokwan Boonyapisit, M.D.
Siriraj Hospital




Which medications?

- ลักษณะการชักและประเภทของโรคลมชักของผู้ป่วย
- การบริหารยา
- ผลข้างเคียงของยากันชัก
- Drug interaction กรณีที่ผู้ป่วยได้ยาหลายชนิดพร้อมกัน
- Special situations
 - Reproductive age
 - Elderly
 - Hepatic impairment
 - Renal impairment



Antiepileptic drug development

Antiepileptic drugs



Special situations



Special situations

- ❖ **Hepatic and renal dysfunction**
- ❖ **Other medical conditions**
 - Transplant patients
 - HIV infected patients
 - Patients with brain tumor
- ❖ **Psychiatric patients**
- ❖ **Elderly**
- ❖ **Women**

Hepatic and renal dysfunction

Hepatic dysfunction



Factors affecting hepatic clearance

- ❖ **The extent of drug binding to the blood component**
- ❖ **Hepatic blood flow**
- ❖ **Hepatic metabolic activity**

AED	Protein binding %	T/2	Site of elimination	Remarks
Gabapentin	0	4-6	Renal, 100% Not metabolize	Dose dependent absorption
Lamotrigine	55	15-30	Hepatic, 90% Glucoronidation	Clearance increased by enzyme inducing AEDs, reduced by VPA
Topiramate	9-17	15-23	Renal, 40-70%	Fraction hepatically metabolized, increased by enzyme inducing AEDs
Levetiracetam	0	6-8	Renal, 66%; hydrolysis of acetamide gr, 34%	Metabolism is nonhepatic hydrolysis
Oxcarbazepine	40	4-9	Hepatic, 70% Hepatic conversion to active metabolite	Based upon 10 Hydroxy carbazepine (MHD), the major active metabolite
Zonisamide	40-60	24-60	Hepatic, 70%	Clearance increased by enzyme inducing AEDs
Pregabalin	0	6	Renal Not metabolize	



Effects	Older AEDs	New AEDs
Measurable increased in free fraction with hypoalbuminemia	PHT VPA	-
Metabolism affected by renal disease	PB	GBP, LEV, TPM
Metabolism affected by liver disease	CBZ, PHT, VPA	LTG, ZNS, OXC, TGB



Dosing adjustment for patients with impaired hepatic function

- ❖ There is insufficient information available to make recommendations on the necessity of dosage adjustment**



Patients with impaired hepatic function

- ❖ Free fractions of **diazepam, PHT, and VPA** increase as a result of reduced circulating albumin concentrations. Frequent serum determinations of free fractions and gradual dose regulations are required.

Renal dysfunction



Effects	Older AEDs	New AEDs
Measurable increased in free fraction with hypoalbuminemia	PHT VPA	-
Metabolism affected by renal disease	PB	GBP, LEV, TPM
Metabolism affected by liver disease	CBZ, PHT, VPA	LTG, ZNS, OXC, TGB



Dosing adjustment for patients with impaired renal function

Creatinine clearance (mL/min)	Dosage (mg)
Gabapentin	
>60	400 tid
30-60	300 bid
15-30	300 od
<15	300 every other day
hemodialysis	200-300* supplement
Levetiracetam	
>80	500-1500 bid
50-80	500-1000 bid
30-50	250-750 bid
<30	250-500 bid
hemodialysis	500-1000*q 24 hr then 250-500 mg supplement
*with supplement dose after HD	



Dosing adjustment for patients with impaired renal function

Creatinine clearance (mL/min)	Dosage (mg)
Topiramate	
>70	Normal dosage
10-70	Decrease dosage 50%
<10	Decrease dosage 75%
hemodialysis	Consider supplement



AEDs that can cause renal stone

❖ **Topiramate**

❖ **Zonisamide**

Using AEDs in patients with other medical conditions



Metabolic pathways of AEDs

CYP 1A2	CYP 2C9	CYP 2C19	CYP 3A4
Carbamazepine*	Phenytoin	Phenytoin*	Carbamazepine
	Phenobarbital	Diazepam	Tiagabine
	Valproate*		Zonisamide
			Ethosuximide
			Felbamate

*Minor metabolic pathway.



Effects on hepatic enzymes

Enzyme inhibitor	Enzyme inducer
Sodium valproate	Phenytoin
	Carbamazepine
	Phenobarbital



Interaction with other drugs

- ❖ **Interaction between CYP3A4 inhibitors and carbamazepine**
- ❖ **Warfarin**
- ❖ **OCPs**
- ❖ **Psychiatric drugs**
- ❖ **Cardiac drugs**
- ❖ **Chemotherapy and immunosuppressive agents**



Commonly used medications that inhibit the CYP3A4

Erythromycin

Clarithromycin

Troleandomycin

Cimetidine

Diltiazem

Verapamil

Fluconazole

Itraconazole

Ketoconazole

Fluvoxamine

Nefazodone

Sertraline

Ritonavir

Indinavir

Nelfinavir

Omeprazole

Propoxyphene



Drug interaction with warfarin

- ❖ **Metabolites through CYP3A4, 2C9**
- ❖ **Phenytoin, phenobarbital and carbamazepine** reduce the concentration of warfarin by up to 50-65%
- ❖ **Phenobarbital and carbamazepine also** reduce the anticoagulation effects of warfarin metabolites
- ❖ **Newer AEDs do not have significant** interaction with anticoagulant



Contents lists available at www.sciencedirect.com

Epilepsy Research

journal homepage: www.elsevier.com/locate/epilepsyres



Short communication

Interactions between non-vitamin K oral anticoagulants and antiepileptic drugs



Claudia Stöllberger*, Josef Finsterer

Krankenanstalt Rudolfstiftung, Wien, Austria

ARTICLE INFO

Article history:

Received 12 April 2016

Received in revised form 1 June 2016

Accepted 24 June 2016

Available online 9 July 2016

Keywords:

Atrial fibrillation

Stroke

Seizure

Drug-drug interaction

ABSTRACT

Atrial fibrillation (AF) is a frequent cause of stroke. Secondary prophylaxis by oral anticoagulants (OAC) is recommended after stroke in AF-patients. OAC can be achieved by vitamin-K antagonists (VKAs) or non-vitamin K antagonist oral anticoagulants (NOACs) like dabigatran, rivaroxaban, apixaban or edoxaban. Seizures are frequent after stroke, and antiepileptic drugs (AEDs) are indicated. The review, based on a literature research, aims to give an overview about pharmacokinetic knowledge and clinical data about drug–drug interactions (DDIs) between NOACs and AED.

Carbamazepine, levetiracetam, phenobarbital, phenytoin and valproic acid might decrease the effect of NOACs by inducing P-glycoprotein (P-gp) activity. Carbamazepine, oxcarbazepine, phenytoin, phenobarbital and topiramate might decrease the effect of NOACs by inducing CYP3A4 activity. Controversial data – inhibition as well as induction of CYP3A4 – were found about valproic acid.

The relevance of these DDIs is largely unknown since there are only sporadic case reports available. To increase the knowledge about DDIs between NOACs and AEDs we suggest subgroup analyses addressing effects and safety of VKAs versus NOACs in patients with AF on AEDs, in case they have been included in previously completed or still ongoing trials or registries. This could be easily feasible and would be desirable in view of the large data already accumulated.

© 2016 Elsevier B.V. All rights reserved.



- ❖ **Intestinal absorption and renal elimination of NOACs are dependent on the intestinal and renal permeability glycoprotein (P-gp) efflux transporter protein system**
- ❖ **Some NOACs are substrates of the hepatic CYP3A4 enzymes**
- ❖ **Induction of P-gp or CYP3A4 might decrease serum NOAC levels, reduce anticoagulant effects and lead to an increase in embolic risk.**



NOAC	P-GP substrate	CYP 3A4 substrate
Dabigatran	Yes	No
Rivaroxaban	Yes	Yes
Apixaban	Yes	Yes
Edoxaban	Yes	Yes

Pharmacokinetic drug interactions mediated by P-gp alone (dabigatran) or in combination with CYP3A4 enzymes (rivaroxaban and apixaban) have been reported

Chin PK, Wright DF, Zhang M, et al. C. Drugs R. D. 2014;14: 113–23.
Hellwig T, Gulseth M. Ann. Pharmacother 2013;47: 1478–87.
Serra W, Li Calz M, Coruzzi P. Clin. Pract 2015; 5: 788



Drug interaction with OCPs

- ❖ **AEDs that cause induction of CYP 3A4 increase metabolism of oral contraceptives resulting in failure of contraceptives.**
- ❖ **Potent enzyme inducing AEDs:**
 - phenytoin, carbamazepine, primidone, phenobarbital.
- ❖ **Less-potent enzyme inducing AEDs:**
 - oxcarbazepine, lamotrigine
 - topiramate >200 mg.

Drug Class

Interactions with AEDs

Antiarrhythmics	Inductor AEDs enhances antiarrhythmics metabolism; phenytoin decreases amiodarone metabolism.
Hypotensive agents	Inductor AEDs enhances beta-blockers and calcium-antagonist metabolism; verapamil and diltiazem inhibit carbamazepine metabolism.
Digoxin	Phenytoin increases digoxin metabolism.
Lipid-lowering drugs	Inductor AEDs enhance lipid-lowering agents metabolism.
Immunosuppressants	Phenytoin, carbamazepine, and barbiturates enhance tacrolimus, sirolimus, and methylprednisolone metabolism.
Antivirals	Inductor AEDs enhance anti-HIV agents metabolism; anti-HIV agents increase carbamazepine, gabapentin, levetiracetam, and lamotrigine levels.
Antibiotics	Carbapenems decrease valproate levels; macrolides increase carbamazepine levels.
Antifungal	Antifungals enhance carbamazepine and phenytoin levels.
Tuberculostatics	Rifampicin enhances phenytoin, carbamazepine, valproate, ethosuximide, and lamotrigine metabolism; isoniazide inhibits it.

Transplant patients



Using AEDs in transplant patients

- ❖ CBZ, oxcarbazepine, PB, and PHT may reduce **cyclosporine, tacrolimus**, and **corticosteroid** blood levels with a delayed effect of up to 10 days.
- ❖ Azathioprine, mycophenolate mofetil, and OKT3 metabolism are not significantly affected by AEDs.

HIV patients



Interaction between ARVs and AEDs

ARV	Protein binding (%)	Metabolism	Potential drugs that may have interaction with AEDs	AEDs that may have interaction with
NRTI	Min- 38	Gluc	↑Zidovudine	VPA
NNRTI	50-90	CYP450		
PI	>99	CYP450	↓Lopinavir/ Ritonavir	PHT

SPECIAL REPORT

Antiepileptic drug selection for people with HIV/AIDS: Evidence-based guidelines from the ILAE and AAN

***†Gretchen L. Birbeck, ‡Jacqueline A. French, §Emilio Perucca, ¶David M. Simpson,
#Henry Fraimow, **Jomy M. George, ††Jason F. Okulicz, ‡‡David B. Clifford,
§§Houda Hachad, and §§René H. Levy for the Quality Standards subcommittee of the American
Academy of Neurology and the ad hoc task force of the Commission on Therapeutic Strategies of
the International League Against Epilepsy**

Epilepsia, 53(1):207–214, 2012



Recommendations

- ❖ AED–ARV administration may be indicated in up to 55% of people taking ARVs.
- ❖ Patients receiving **phenytoin** may require a **lopinavir/ritonavir (PI)** dosage increase of approximately 50% to maintain unchanged serum concentrations (Level C: one class II study).
- ❖ Patients receiving **valproic acid** may require a **zidovudine (NRTI)** dosage reduction to maintain unchanged serum zidovudine concentrations (Level C).
- ❖ Coadministration of valproic acid and efavirenz (NNRTI) may not require efavirenz dosage adjustment (Level C: one class II study).



Recommendations

- ❖ It may be important to avoid **enzyme inducing AEDs** in people on ARV regimens that include **protease inhibitors or non nucleoside reverse transcriptase inhibitors** because pharmacokinetic interactions may result in virologic failure, which has clinical implications for disease progression and development of ARV resistance. If such regimens are required for seizure control, patients may be monitored through pharmacokinetic assessments to ensure efficacy of the ARV regimen (Level C: one class II study).

Brain tumors



Potentials interaction between AEDs and chemotherapy

- ❖ Enzyme inducing AEDs have been shown to have effects on levels of chemotherapy that metabolite through CYP 450
- ❖ Taxanes, vinca alkaloids, methotrexate, teniposide, and camptothecin analogues such as irinotecan

Vecht CJ, Wagner GL, Wilms EB. Lancet Neurol 2003;2:404–9.



Potentials interaction between AEDs and chemotherapy

- ❖ In a study of 716 children with ALL, 40 children who were on enzyme-inducing AEDs had worse event-free survival (hazard ratio 2.67 [95% CI, 1.50 to 4.76]), hematological relapse (3.40 [1.69 to 6.88]) and CNS relapse (2.90 [1.01 to 8.28]).
- ❖ These children were found to have a higher clearance of teniposide and methotrexate.

Relling MV, Pui CH, Sandlund JT, et al. Lancet 2000;356:285–90



Potentials interaction between AEDs and chemotherapy

- ❖ In a study on glioblastoma multiforme treated with adjuvant CCNU after surgery and radiotherapy, patients receiving enzyme-inducing AEDs (carbamazepine in 80% of patients) had a significantly shorter survival, 10.8 versus 13.9 months, than patients treated with non-enzyme-inducing AEDs (valproic acid in 80% of patients)

Oberndorfer S, et al. J Neurooncol 2005;72:255–60



Patients with brain tumors

- ❖ **Enzyme-inducing AEDS can interfere with the level of concomittent chemotherapy and should be avoided.**
- ❖ **Valproic acid may be considered as a first-line agent, although physicians should be aware of the potentially enhanced toxicity of concomitant agents that share the same P-450 coenzyme metabolic pathway.**



Patients with brain tumors

- ❖ **Newer AEDs that do not metabolite through CYP 450 system also can be used.**
- ❖ **More evidence is still needed.**



www.thaiepilepsysociety.com



สมาคมโรคลมชักแห่งประเทศไทย Thai Epilepsy Society

[Home](#) [เกี่ยวกับสมาคม](#) [ข่าวประชุมวิชาการ](#) [สไลด์การประชุมวิชาการ](#) [ความรู้สำหรับแพทย์](#) [ความรู้สำหรับประชาชน](#) [Links](#) [ติดต่อสมาคม](#)

การประชุมวิชาการประจำปี 2556



การประชุมวิชาการประจำปี 2556 สมาคมโรคลมชักแห่งประเทศไทย จัดขึ้นระหว่างวันที่ 25-26 กรกฎาคม 2556 ณ โรงแรมสุโกศล กรุงเทพฯ สามารถดูรายละเอียดของการประชุมได้ที่ข่าวประชุมวิชาการ

Recent Posts

[การประชุม ASEPA & EST Workshop on Epilepsy Surgery](#)

[การประชุมวิชาการประจำปี 2556](#)

Meta

[Log in](#)

[Entries RSS](#)

[Comments RSS](#)